

XP-002199667

AN - 1985-287885 [25]
AP - JP19840054862 19840322
CPY - KYOW
DC - B02 D16
FS - CPI
IC - C12P17/06 ; C12R1/62
MC - B06-A01 D05-C
M2 - [01] D013 D022 D023 D120 G013 G100 H4 H402 H442 H541 H8 J5 J521 M1
M113 M210 M211 M272 M280 M281 M320 M412 M511 M520 M531 M540 M720 M903
N132 N425 N512 N513 Q233
PA - (KYOW) KYOWA HAKKO KOGYO KK
PN - JP60199396 A 19851008 DW198546 005pp
PR - JP19840054862 19840322
XA - C1985-124869

XIC - C12P-017/06 ; C12R-001/62

AB - J60199396 Fermentative prepn. of daidzein (I) and tectorigenin (II)
comprises culturing microorganisms to produce and isolate (I) and (II)
from the culture fluid.

- (III) can be cultured with usual culture fluid for actinomyces, contg.
suitable C-source, N-source, inorganic salts, etc.. Carbohydrates
(glucose, fructose, sucrose, starch, dextrin, mannose, maltose,
molasses), organic acids (citric acid, malic acid, acetic acid,
fumaric acid), alcohols (methanol, ethanol), hydrocarbons (methane,
ethane, propane, n-paraffine), amino acids, glycerol etc. can be used
as C-source. Ammonium salts (ammonium chloride, ammonium sulphate,
ammonium nitrate, ammonium phosphate), aminoacids (aspartic acid,
glutamine, cystine, alanine) and other nitrogen contg. substances can
be used as N-source. Other substances, accelerating prodn. of (I)
and/or (II) may be added to the culture fluid as necessary.

Cultivation is pref. carried out at 20-40 deg. C, pH 7, for 3-15 days,
by shaking or aerobic culture. Concn. of (I) or (II) may reach a
maximum. (I) and (II) can be isolated from culture fluid by
combination of filtration, adsorption chromatography, thin-layer
chromatography etc..

- USE/ADVANTAGE - (I) and (II) are known and methods to isolate them are
already reported. By cultivation of the new microorganism,
Streptovorticillium sp. K-251 (III), a mixt. of (I) and (II) is obtd.,
and then (I) and (II) are isolated from culture fluid.

IW - FERMENTATION PREPARATION ISO FLAVONE DERIVATIVE DAIDZEIN CULTURE
STREPTOVERTICILLIUM SPECIES

IKW - FERMENTATION PREPARATION ISO FLAVONE DERIVATIVE DAIDZEIN CULTURE
STREPTOVERTICILLIUM SPECIES

NC - 001

OPD - 1984-03-22

ORD - 1985-10-08

PAW - (KYOW) KYOWA HAKKO KOGYO KK

TI - Fermentative prepn. of iso-flavone derivs. daidzein and tectorigenin -
by culturing streptovorticillium Sp. K-251